

PRESSURE INDUCED TOPOLOGICAL PHASE TRANSITIONS IN MEMBRANES

P.T.C. So*, Sol M. Gruner and Shyamsunder Erramilli¹

Department of Physics, Joseph Henry Laboratories Princeton University P.O. Box 708 Princeton, NJ 08544

Some highly unusual features of a lipid-water liquid crystal are revealed by high pressure x-ray diffraction, light scattering and dilatometric studies of the lamellar (bilayer L_α) to nonlamellar inverse hexagonal (H_{II}) phase transition. (i) The size of the unit cell of the H_{II} phase increases with increasing pressure. (ii) The transition volume, ΔV_{bh} , decreases and appears to vanish as the pressure is increased. (iii) The intensity of scattered light increases as ΔV_{bh} decreases. Data are presented which suggest that this increase is due to the formation of an intermediate cubic phase, as predicted by recent theoretical suggestions of the underlying universal phase sequence.

PACS numbers: 64.70.Md, 87.22.Bt

Dispersions of biological lipids in water exhibit a rich variety of structurally complex phases as the temperature, pressure and water content are varied. The lamellar liquid crystalline bilayer (L_α) to inverse hexagonal (H_{II}) phase transition has attracted particular interest because it is the simplest of the lyotropic transitions in which both the topology and curvature of the lipid-water interfaces internal to the liquid crystal change discontinuously at the transition [2].

The presence of water and electrostatic effects makes lyotropic dispersions very complicated and difficult to understand. Insight into understanding such systems first came with the recognition that they form assemblies of 2-dimensional liquid-like films which elastically resist lateral extension or compression, or bending away from a desired curvature. Winsor [3] and Helfrich [4] suggested that much of the complexity of the molecular interactions can be lumped into phenomenological parameters such as the curvature of the lipid-water interfaces and the elastic moduli of the lipid layers. The utility of this phenomenological approach toward understanding the lamellar bilayer to nonlamellar transitions has been established by Gruner and coworkers [5] and by Charvolin [6].

The idea that many aspects of the global phase behavior can be explained without considering the detailed molecular structure of the molecules involved led Goldstein and Leibler [7] to propose a mean-field model for the well known bilayer chain melt (i.e. gel-liquid crystalline) phase transition. They proposed a Ginzburg-Landau approach using the strain field of the lipid layers, with phenomenological parameters modulated by microscopic interactions arising from van der Waals, hydration and electrostatic forces. While this approach has not been extended to nonlamellar phases, the enormous simplification afforded by Helfrich's insight has stimulated recent experimental [2] and theoretical interest in nonlamellar phases [8,9]. A rather general attempt in this direction is the Landau- Ginzburg model of Chen et al [9], which involves a combination of scalar and vector order parameters that is claimed to have, in principle, sufficient complexity to describe not only lamellar and hexagonal phases, but also more geometrically complex phases, such as bicontinuous cubic phases [8,10]. However, the large number of phenomenological parameters involved, as well as a lack

of an intuitive physical interpretation or ready experimental handle on the order parameters, suggests that deeper insight may be derived from simpler explanations of unusual phase or structural behavior in cases where care has been taken to limit the energetic significance of complicating factors, such as the effects of electrostatic charge or limited water. Although there have been numerous studies of the effects of variation of temperature on these transitions [2], there have been very few investigations of the effects of pressure. In this Letter, we describe some rather unusual phenomena exhibited by excess-water dispersions of electrically neutral lipid, 1,2-dioleoyl-sn-glycero-3-phosphoethanolamine (DOPE; also known as dioleoylphosphatidylethanolamine), which swells in water to form phases with a well-defined water fraction, with a view to stimulating quantitative models.

1) *Nonlamellar phase transitions temperatures are very pressure sensitive.* Optical scattering [11] and x-ray diffraction [12] measurements suggested that the L_α - H_{II} transition temperature may be depressed by surprisingly modest pressures. Our experiments were conducted using a rotating anode x-ray source, a point-focussed beamline, a thermostated high pressure beryllium x-ray cell, and a two-dimensional x-ray detector [13] which, in combination, allowed rapid surveys of the temperature-pressure structure of small (~ 10 mg) lipid samples. The phases and internal dimensions of the lipid-water liquid crystals were readily identified by the characteristic x-ray diffraction patterns of unoriented dispersions [14]. The Clausius-Clapeyron coefficient of change of transition temperature with pressure was $dT_{bh}/dP = 44$ K/kbar, a value which is nearly twice as large as for the chain melting transition [15] and which establishes the excess-water L_α - H_{II} transition as one of the most pressure sensitive first order transitions known in a non-gaseous system. The isothermal bulk compressibility of the lipid-water system is comparable to that of water (see below). Since the system is not unusually compressible, this suggests that the L_α - H_{II} phase transition involves a very delicate balance of competing energetic contributions such that small changes in overall volume result in large changes in the balance.

2) The unit cell spacing compressibility is negative. Figure 1 shows that the unit cell spacing, b , of the H_{II} phase increases as pressure is applied. Furthermore, this pressure

dependence is remarkably steep: at 20 C, b increases by nearly 1 Å for every 100 bar increase in pressure. The increase in the unit cell is almost all due to the transfer of water from the coexisting water bulk to the liquid crystal [16]. The data yield a value of the isothermal bulk linear compressibility of the Bragg lattice of $\alpha_T = -\frac{1}{b} \left(\frac{\partial b}{\partial P} \right)_T = -2.5 \times 10^{-2} \text{ bar}^{-1}$. Not only is α_T negative, but it is more than 1500 times as large as the compressibility of water. We define a generalized Clausius-Clapeyron relation, $\frac{dT}{dP} = \frac{\Delta T}{\Delta \xi} \frac{\Delta \xi}{\Delta P}$, that relates the sensitivity of a structural parameter, ξ , to pressure and temperature changes. We find for the unit cell spacing with $\xi = b$, $\Delta b/\Delta T = 0.22 \text{ Å/K}$ and $\Delta b/\Delta P = 9.6 \text{ Å/kbar}$, giving for $\frac{dT}{dP} = 44 \text{ K/bar}$, which is essentially the same as the Clausius-Clapeyron value for the L_α - H_{II} phase transition given above. *This suggests that the thermodynamics of the phase transition may be understood using models based only on the gross structural parameters, without reference to details of the molecular interactions.*

Thermodynamic stability arguments show that the apparent negative compressibility in Figure 1 refers only to the increase in the unit cell size of the two-dimensional hexagonal lattice; the overall specimen volume must decrease with increasing pressure. Almost all of the increase in size with increasing pressure arises from an increase in the radius of the central water core of the hexagonally packed tubes and the resultant influx of bulk water. Electron density reconstruction of the x-ray data show that the truly pressure sensitive aspect of the system is the spontaneous curvature of the lipid layer making up the hexagonally packed tubes [16]. Measurement of parameters such as the average volume, length and interfacial area per lipid molecule cannot be obtained from x-ray diffractometry alone; therefore we constructed a high pressure (0 - 3 kbar) dilatometer capable of resolving volume changes of about 0.1 nl/mg in small lipid- water specimens [17]. Figure 2 shows the specific volume as a function of pressure at various temperatures.

3) At least four different pressure sensitive phase transitions are observed in excess water-DOPE samples (Fig.2). The isothermal bulk compressibility of the DOPE-water system is about $\beta_T = -\frac{1}{V} \left(\frac{\partial V}{\partial P} \right)_T = +2 \times 10^{-4} \text{ bar}^{-1}$. Phase transitions are readily identified

in the dilatometry data as discontinuities in the lipid specific volume or as slight changes in the slope of the isotherms signalling a change in compressibility. Figure 2 shows that four distinct transitions corresponding to five phases, which have also been examined via x-ray diffraction [18]. (i) At 13 C a small volume change occurs at about 200 bar which corresponds to the L_α - H_{II} transition. (ii) At 8 C and 750 bar, a large volumetric change is observed which is identified from the appearance of sharp peaks in the wide angle x-ray scattering (WAXS) as the familiar chain melting (gel to liquid crystalline) transition. The volume change is in agreement with previous thermodynamic studies on lipid systems with phosphatidylcholine (PC) headgroups [21]. (iii) At 8 C and 1250 bar, a phase transition with a small volume change is observed with the position and intensity of the sharp peaks in the WAXS changing discontinuously, suggesting that the symmetry of the packing of the hydrocarbon chains has been changed. (iv) A large volume change is observed at 8 C and 2 kbar the system forms a high-pressure phase characterized by five distinct WAXS lines. This suggests that the chains are packed with a high degree of order and that the phase is likely to be a crystalline phase. Crystalline phases corresponding to sub-gel transitions are known in PC systems [20,21].

4) The transition volume, ΔV_{bh} , enthalpy, ΔH_{bh} and entropy, ΔS_{bh} , of the L_α - H_{II} transition decreases as the pressure is increased. The transition volume was measured by dilatometry and the transition enthalpy and entropy were then computed using the Clausius-Clapeyron relation. The transition volume decreases as the temperature increases (inset, Figure 2). Most interestingly, ΔV_{bh} , and therefore ΔH_{bh} and ΔS_{bh} , become too small to measure with our apparatus at about 80 C and ~ 1750 bar.

5) Light scattering of the DOPE-water system increased as the transition volume decreased. A conventional high pressure optical cell was constructed out of beryllium- copper and equipped with sapphire windows and unsupported area seals. The intensity, $I(q)$, of scattered He-Ne laser light ($\lambda = 632$ nm) was measured with a light sensitive diode at an angle corresponding to $q \sim 10^{-3}$ nm $^{-1}$. A plot of $I(q)$ shows a prominent peak at a temperature in agreement with the L_α - H_{II} transition, as identified by both dilatometry and x-ray

diffraction.

6) A cubic phase occurs between the L_α and H_{II} phases. It might be imagined that the increase in intensity of the scattered light concomitant with a decrease in transition volume is suggestive of some type of critical behavior. However, a different hypothesis provides a more plausible explanation: the light-scattering anomalies observed in the L_α - H_{II} transition are due to the onset of an intermediate phase, associated with very little latent heat, but whose formation is kinetically hindered by large activation barriers. Defects forming the precursor to the intermediate phase scatter light.

In order to test this hypothesis, an excess water-DOPE specimen was allowed to equilibrate at 80 ± 0.05 C and 1750 ± 3 bar, the nominal ‘critical point’. In situ x-ray diffraction patterns of the sample were taken roughly every hour for three days. The initial diffraction pattern (Figure 4a) is characteristic of the hexagonal phase. As time elapses, diffuse scatter becomes prominent concomitant with a decrease in the intensity of the hexagonal diffraction peaks. Within 12 hours, additional peaks appear. After 36 hours, the diffraction pattern can be readily indexed to a cubic lattice of $Pn3m/Pn3$ symmetry. The x-ray data shows that a cubic phase forms at high temperature and pressure. The spontaneous formation of the cubic phase at the expense of the hexagonal phase indicates that the cubic phase is stable. The slow formation of the cubic phase and the coexistence seen in Figure 4 with the metastable hexagonal phase are indicative of a pronounced kinetic barrier between the phases.

The sequence of phase dimensionality, from 1-D lamellar to 3-D cubic to 2-D hexagonal, underscores the fact that the lattice dimensionality is not a relevant parameter for these topological phase transitions. Gruner [5], and Charvolin [6] suggest that the phase behaviour can be understood as a result of a *frustration* between opposing requirements of membranes that have to curl to satisfy curvature requirements on the one hand and to try to maintain constant bilayer thickness on the other. Because the phases are formed by a subtle balance of competing interactions, the latent heat associated with the transitions can be small. Anderson et al [8] argued that certain bicontinuous cubic phases provide equilibrium

compromises between lamellar and inverse hexagonal phases such that one expects them to always occur between these phases in the phase diagrams. The data presented in this Letter support this picture, and suggest that the underlying universality of this phase sequence may be obscured by the large kinetic barriers. The origin of the large barriers lies in the fact that the L_α , H_{II} , and bicontinuous cubic phases have very different topologies of the interfaces. Transitions between lamellar and non-lamellar phases necessarily require lipid-water interfaces to be torn and reformed. The process requires exposing the hydrocarbon tails of the lipid molecules to water. The energy associated with this disruption probably is the large activation barrier.

Rather striking support for the notion of an underlying universality comes from the observation diblock copolymers exhibit essentially the same sequence of phases [22], although the microscopic interactions are quite different. It would be interesting to see if the observed negative anisotropic compressibility also carries over to the diblock copolymer systems. More importantly, a unified quantitative theory of all systems exhibiting topological phase transitions may now be possible.

I. ACKNOWLEDGEMENTS

This work was supported by the ONR (Contract N00014-86-K-0396 P0001), and by the DOE (Grant DE-FG02-87ER60522) and by the NIH (Grant GM32614). We thank R. Bruinsma, J. T. Gleeson, D. C. Turner and M. W. Tate for useful discussions and assistance with figures.

REFERENCES

* Present address: Physics dept., University of Illinois, Urbana IL-61801.

- [1] To whom correspondence should be addressed.
- [2] Some recent reviews of non-bilayer phases are: J. M. Seddon (1990), *Biochim. Biophys. Acta***1031**, 1-69; G. Lindblom and L. Rilfors, *Biochim. Biophys. Acta***988**, 221-256 (1989); K. Larsson, *J. Phys. Chem.* **93**, 7304 (1989); M. W. Tate, E. F. Eikenberry, D. C. Turner, E. Shyamsunder and S. M. Gruner (1991), *Chem. Phys. Lipids.* **57**, 147-164.
- [3] P. A. Winsor *Mol. Cryst. Liq. Cryst.* **12**, 141-178 (1971).
- [4] W. Helfrich *Z. Naturforsch A* **28C**, 693 (1973).
- [5] G. L. Kirk, S. M. Gruner and D. L. Stein, *Biochem.* **26**, 6592-6598 (1984).
- [6] J. Charvolin *J. Physique* **46**, C3-173 to C3-190 (1985).
- [7] R. E. Goldstein and S. Leibler, *Phys. Rev. Lett.* **61**, 2213 (1988).
- [8] D. M. Anderson, S. M. Gruner and S. Leibler, *Proc. Natl. Acad. Sci.* (1988).
- [9] K. Chen, C. Jayaprakash, R. Pandit and W. Wenzel, *Phys. Rev. Lett.* **65**, 2736 (1990).
- [10] S. T. Hyde, S. Adersson, B. Ericsson and K. Larsson, *Z. Kristall.* **168**, 213-219 (1984).
- [11] P. Yeager and E. L. Chang, *Biochimica. et. Biophysica. Acta* **731**, 491-494 (1983).
- [12] E. Shyamsunder, P. Botos and S.M. Gruner, *J. Chem. Phys.* **90**, 1293-1295 (1989).
- [13] Details of the apparatus can be found in P. T. C. So, S. M. Gruner and E. Shyamsunder, *Rev. Sci. Instrum.* **63** 1763-1770 (1992).
- [14] V. Luzzati in *Biological Membranes*, Vol. I, (D. Chapman, Ed.), Academic, New York pp 71-123 (1968).
- [15] J. Stamatoff, D. Guillam, L. Powers, P. Cladis and D. Aadsen (1978), *Biochem. and*

Biophys. Res. Comm. **85**, 724-728.

- [16] O. Narayan, P.T.C. So, D.C. Turner, S.M. Gruner, M.W. Tate and E. Shyamsunder, Phys. Rev. A **42**, 7479-7482 (1990).
- [17] P. T. C. So, S. M. Gruner and E. Shyamsunder Rev. Sci. Instr. (in press).
- [18] P. T. C. So, Ph. D. Thesis, Princeton University. The work reported here forms a part of this thesis.
- [19] E. Shyamsunder, S.M. Gruner, M.W. Tate, D.C. Turner, P.T.C. So and C.P.S. Tilcock (1988), Biochem. **27**, 2332.
- [20] P. T. T. Wong Ann. Rev. Biophys. Bioeng. **13**, 1-24 (1984).
- [21] Utoh and Takumura (1985), Japanese Journal of Applied Physics *24*, 356-360.
- [22] S.M. Gruner, J. Phys. Chem. **93**, 7562-7570 (1989).
- [23] D. A. Wilkinson and J. F. Nagle, Analytical Biochemistry **84**, 263-271 (1978).

FIGURES

FIG. 1. *Unit cell spacing in the H_{II} phase as a function of pressure at various temperatures.*

Data are shown for aqueous dispersions of high purity DOPE purchased from Avanti Polar Lipids at the temperatures 25 C ($\circ$), 43 C (∇), 62 C ($\triangle$) and 81 C ($\square$). Sample purity was periodically checked before and after experiments by thin layer chromatography (TLC). Lyophilized lipid was mechanically mixed with excess deionized water and thermally cycled several times between -10°C and 20°C to ensure mixing. The hydrated lipid was transferred to a high-pressure cell designed for x-ray measurements. The dashed line indicates the approximate phase boundary between the H_{II} and L_α phases.

FIG. 2. *$P - V$ isotherms for the DOPE-water system.*

Solid lines indicate experimental data. Isotherms at 5⁰ C intervals from 8 to 83 ⁰C (from left to right) are shown. The phase transitions indicated are (i) the inverted hexagonal to lamellar transition $H_{II} - L_\alpha$; (ii) the chain melting transition $L_\alpha - L_\beta$; (iii) and (iv) are gel-gel phase transitions not previously reported in the DOPE-water system, but are presumed to be similar to reported transitions in other lipid-water systems [20]. Inset shows that the transition volume ΔV_{bh} for the $L_\alpha - H_{II}$ transition derived from the isotherms using a procedure similar to that of Wilkinson and Nagle [23]. ΔV_{bh} decreases as function of pressure and becomes too small to measure as the pressure reaches ~ 1750 bar.

FIG. 3. *Light scattering study of the DOPE-water system.*

Data were taken at a fixed value of the magnitude of the scattering vector ($q = 10^{-3}$ nm⁻¹), at the pressures indicated. The increase in light scattering intensity coincides with the decrease in ΔV_{bh} .

FIG. 4. *X-ray diffraction studies at the point at which ΔV_{bh} becomes too small to measure.*

Evolution of the x-ray diffraction pattern at the times indicated after reaching 1750 bar, 80 C. The locations of the Bragg peaks in the diffraction pattern is characteristic of an inverted hexagonal phase (indexed in curly brackets). As time elapses, additional Bragg peaks emerge that index to a cubic lattice (square brackets). The best fit to the observed Bragg peaks corresponds to a lattice of $Pn3m$ (or $Pn3$) symmetry shown schematically in the lower left. that has been observed in other lipid-water systems. The cubic phase may also be obtained by repeatedly thermally cycling the system across the $L_\alpha - H_{II}$ phase transition [19].